



**Figure 3.4-38.** Year-round and Seasonal Trawl Restrictions in Prince William Sound




Source: NMFS 2001b

**Figure 3.4-39.** State Groundfish Regions




Source: ADF&G 2002c

**Figure 3.4-40.** Alaskan Herring Fisheries




Source: ADF&G 2002a

**Figure 3.4-41.** Herring Sac Roe Harvests, 1963-2001




Source: ADF&G 2001

**Figure 3.4-42.** Drift Gillnet




Source: K. Richards, Tetra Tech FW, Bothell, WA


**Figure 3.4-43.** Purse Seine Gear




Source: Blackwell Science (Jennings et al. 2001)  
Permission to use figure requested by Craig Rose (pending)




**Figure 4.3-1.** Blue king crab female abundance from trawl survey data and blue king crab catch data, by State of Alaska statistical area.




**Figure 4.3-2.** Red king crab female abundance from trawl survey data and red king crab catch data, by State of Alaska statistical area.



**Figure 4.3-3.** Snow crab female abundance from trawl survey data and snow crab catch data, by State of Alaska statistical area.



**Figure 4.3-4.** Tanner crab female abundance from trawl survey data and Tanner crab catch data, by State of Alaska statistical area.



**Figure 4.3-5.** Locations of commercial scallop beds in Alaska.